



Published in final edited form as:

Depress Anxiety. 2011 August ; 28(8): 648–657. doi:10.1002/da.20865.

Complicated grief associated with Hurricane Katrina

M. Katherine Shear, MD, Katie A. McLaughlin, PhD, Angela Ghesquiere, MSW, Michael J. Gruber, MS, Nancy A. Sampson, BA, and Ronald C. Kessler, PhD

Columbia University School of Social Work and Columbia University College of Physicians and Surgeons, New York (Drs. Shear & Ms. Ghesquiere); Division of General Pediatrics, Children's Hospital Boston & Harvard Medical School (Dr. McLaughlin), Department of Health Care Policy, Harvard Medical School, Boston (Mr. Gruber, Mrs. Sampson, and Dr. Kessler)

Abstract

Background—Although losses are important consequences of disasters, few epidemiological studies of disasters have assessed complicated grief (CG) and none assessed CG associated with losses other than death of loved one.

Methods—Data come from the baseline survey of the Hurricane Katrina Community Advisory Group (CAG), a representative sample of 3,088 residents of the areas directly affected by Hurricane Katrina. A brief screen for CG was included containing four items consistent with the proposed DSM 5 criteria for a diagnosis of bereavement-related adjustment disorder.

Results—58.5% of respondents reported a significant hurricane-related loss: Most-severe losses were 29.0% tangible, 9.5% interpersonal, 8.1% intangible, 4.2% work-financial, and 3.7% death of loved one. 26.1% of respondents with significant loss had possible CG and 7.0% moderate-severe CG. Death of loved one was associated with the highest conditional probability of moderate-severe CG (18.5%, compared to 1.1–10.5% conditional probabilities for other losses) but accounted for only 16.5% of moderate-severe CG due to its comparatively low prevalence. Most moderate-severe CG was due to tangible (52.9%) or interpersonal (24.0%) losses. Significant predictors of CG were mostly unique to either bereavement (racial-ethnic minority status, social support) or other losses (pre-hurricane history of psychopathology, social competence.).

Conclusions—Non-bereavement losses accounted for the vast majority of hurricane-related possible CG despite risk of CG being much higher in response to bereavement than to other losses. This result argues for expansion of research on CG beyond bereavement and alerts clinicians to the need to address post-disaster grief associated with a wide range of losses.

Keywords

bereavement; loss; grief; disaster mental health; complicated grief; loss; suicide

Correspondence: M. Katherine Shear, M.D., Columbia University School of Social Work, 1255 Amsterdam Avenue, New York, NY 10027. ks2394@columbia.edu; Phone: (212) 851-2176; Fax: (212) 851-2175.

Data access and responsibility: Dr. Kessler had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Financial Disclosures: Dr. Kessler has been a consultant for GlaxoSmithKline Inc., Kaiser Permanente, Pfizer Inc., Sanofi-Aventis, Shire Pharmaceuticals, and Wyeth-Ayerst; has served on advisory boards for Eli Lilly & Company and Wyeth-Ayerst; and has had research support for his epidemiological studies from Bristol-Myers Squibb, Eli Lilly & Company, GlaxoSmithKline, Johnson & Johnson Pharmaceuticals, Ortho-McNeil Pharmaceuticals Inc., Pfizer Inc., and Sanofi-Aventis. The remaining authors report nothing to disclose.

Death of a loved one is one of the many types of loss caused by natural disasters.^[1–16] Until very recently, though, loss was evaluated in epidemiological studies of the psychiatric consequences of disaster only as a stressor that might trigger PTSD or depression.^[10,11] There is now growing awareness of the clinical significance of complicated grief (CG)^[17] with recognition of its potential importance following disasters.^[18–20] Uncomplicated acute grief is often intense and disruptive shortly after the occurrence of a loss^[21] but typically becomes more muted and less impairing over time.^[22] For people with CG, though, this transformation does not occur and acute grief symptoms (e.g., strong feelings of longing-yearning, preoccupation with thoughts-memories of the deceased, withdrawal, loss of interest) persist longer. Research has documented that CG is impairing,^[23–25] distinguishable from major depression and post-traumatic stress disorder,^[23–27] and profits from psychotherapy that focuses on complicated grief symptoms rather than on depression^[28,29] or on general support.^[30]

A DSM 5 workgroup has proposed that CG be added to DSM 5 as a new diagnosis of bereavement-related adjustment disorder, which is described as including intense yearning, difficulty accepting, and anger over the death of a loved one along with a feeling that life is empty or meaningless.^[31] Although this proposal stipulates that these symptoms occur following the loss of a loved one, evidence exists that grief symptoms also often occur after non-bereavement losses, such as job loss,^[32] loss of a home,^[33] loss of the ability to function,^[34,35] receiving a diagnosis of a fatal disease,^[36–38] and caring for a loved one with dementia.^[39,40] Grief symptoms associated with these non-bereavement losses appear to be similar to those associated with bereavement.^[35] Yet we are aware of no disaster research on CG associated with these non-bereavement losses.

The current report presents preliminary data on this issue from a brief screening scale of CG collected in a survey with the Hurricane Katrina Community Advisory Group (CAG), a representative sample of pre-hurricane residents of the areas in Alabama, Louisiana, and Mississippi directly affected by Hurricane Katrina.^[41–44] Hurricane Katrina was one of the most devastating natural disasters in U.S. history^[45] and was associated with substantial losses of life, property, income, and community,^[41,46] providing an excellent opportunity to examine the prevalence and correlates of CG associated with a range of disaster-related losses.

METHODS

Sample

The CAG is a representative sample of 3,088 English-speaking adults (≥ 18 years old) recruited from random-digit-dial telephone calls of households in the FEMA-defined disaster area affected by Hurricane Katrina and from random selection of families applying for American Red Cross assistance. The baseline CAG interviews, the focus of the current report, were carried out with separate samples of respondents in three waves: January-March 2006 (5–7 months post-hurricane) $n = 1,043$; April-June, 2006 (7–10 months post-hurricane) $n = 723$; December, 2006-April, 2007 (15–19 months post-hurricane) $n = 1,322$. Interviews were carried out in these three waves based on added funding that allowed baseline sample size to increase on two different occasions after the first wave. The 3,088 total respondents had a low (35.2%) cooperation rate (i.e., the survey completion rate among pre-designated respondents who were successfully traced to their current residence at the time of interview) due to the requirement that respondents make a long-term commitment to remain in the CAG (and provide contact information for an informant who would know their whereabouts if they moved), as we aimed to use the CAG to track the progress of post-hurricane recovery over time.

A non-response survey found that CAG non-respondents were similar to respondents on socio-demographic variables, but had somewhat higher hurricane-related stress on a 0–10 scale (where 0 meant “no stress at all” and 10 meant “the most stress you can imagine”) and more psychological distress (assessed with a short series of questions about common anxiety-mood symptoms scored on a 0–10 scale) than respondents. The median and interquartile range (IQR: 25th–75th percentiles) of hurricane-related stress were 8.0 (6.0–10.0) among non-respondents and 7.0 (5.0–9.0) among respondents. The median and IQR of psychological distress were 2.9 (1.2–4.4) among non-respondents and 1.7 (0.6–3.5) among respondents. A weight was applied to the baseline CAG data to adjust for these response biases. No data were collected from non-respondents, though, on specific hurricane-related losses. A within-household probability of selection weight and post-stratification weight were also used to adjust for residual discrepancies between the CAG and the 2000 Census population on a range of socio-demographic and pre-hurricane housing variables. The consolidated CAG sample weight was then trimmed to increase design efficiency.

Measures

Hurricane-related stressors—Respondents were asked 30 questions regarding exposure to hurricane-related stressors. Categories of stressors sufficiently common to be analyzed in subgroup analyses were serious risk of death, death of a family member or close friend, victimization due to lawlessness after the storm (e.g., robbery or physical assault), victimization of a loved one, physical illness or injury caused or exacerbated by the storm, extreme physical adversity (e.g., sleeping in a church basement, difficulty obtaining food or clothing), extreme psychological adversity (e.g., living in distressing circumstances, such as having to use the toilet or change clothes without adequate privacy), major property or income loss, and ongoing difficulties associated with housing (e.g., experiencing multiple moves or living in substantially worse post-hurricane than pre-hurricane housing).^[41]

Loss events—Respondents were asked to nominate and rate their one “most significant hurricane-related loss” on a 0–10 scale (where 0 meant “no loss” and 10 meant “the greatest loss you can imagine”). Only respondents who rated their loss 3+ were administered the grief questions described below. These losses were coded as death of loved one, work/financial losses, other tangible losses (e.g., home, possessions, memorabilia), intangible losses (e.g., quality of life, sense of well-being, control, security, way of life), and interpersonal losses (e.g., separations from family or friends, reduced quality of relationships with family or friends). A small proportion of respondents did not respond to this question (1.2%) or provided uncodable responses (2.7%).

Grief—Respondents with losses rated 3+ were asked four questions about grief in the past 30 days associated with their most significant loss: (1) *How often have you found yourself longing or yearning for the people or things you lost?* (2) *How bitter do you feel over your loss?* (3) *How empty or meaningless do things seem since your loss?* and (4) *How difficult is it for you to accept your loss or to believe that it’s real?* Response options were coded 0–4 (*almost all, most, some, a little, and none of the time* for the first question; and *not at all, a little, some, a lot, and extremely* for the other three questions). Principal axis factor analysis found one meaningful factor with an unrotated eigenvalue of 2.7 compared to 0.5 for the second factor and factor loadings of .77+ for all items. A composite measure of grief was created by reflecting responses to the first question and then summing the four responses to create a 0–16 scale. A preliminary categorical classification was made with the following categories: *severe CG* (15–16), *moderate CG* (13–14), *mild CG* (8–12), *sub-threshold grief* (5–7), and *no- minimal grief* (0–4). These cut-points were based on an analysis of sensitivity-specificity of responses to a similar subset of questions in reproducing diagnoses of CG based a widely utilized CG scale^[47] in a large clinical dataset.^[48] The fact that these

interviews took place 5–19 months after the hurricane suggests that the majority of cases had symptoms for more than six months, which is the minimum duration typically specified for CG.^[49,50]

Socio-demographics—We examined associations of CG symptoms with age, sex, race/ethnicity (Non-Hispanic White versus Other), education, marital status, family income in the year before the hurricane (low/low-middle vs. high-middle/high), health insurance status, and pre-hurricane location of residence (New Orleans Metropolitan Area versus elsewhere). Low/low-middle family income was defined as less than or equal to the population median on the ratio of pre-tax income to number of family members, whereas high/middle-high income was defined as greater than the median on this ratio.

Mental illness—Respondents completed short screening scales of pre-hurricane lifetime history of major depressive episode, anxiety disorders (panic disorder, generalized anxiety disorder, post-traumatic stress disorder), substance use disorders (alcohol or drug abuse with or without dependence), intermittent explosive disorder, and suicidality. These scales were adapted from the Family History Research Diagnostic Criteria interview^[51] and its extensions.^[52] The questions about suicidality were taken from the Self-Injurious Thoughts and Behaviors Interview.^[53] The K6 scale of nonspecific psychological distress^[54] was used to screen for DSM-IV Serious Mental Illness (SMI) in the 30 days before interview. Validation studies have found area under the receiver operating characteristic curve of 0.86–0.89 of the K6 predicting clinical diagnoses of SMI.^[54–56] K6 scores of 13–24 were classified probable SMI. A small K6 clinical reappraisal study (n=15) in the CAG selected 8 respondents with K6 scores in the clinical range and 7 below that range and blindly administered the Structured Clinical Interview for DSM-IV (SCID).^[57] Sensitivity and specificity were perfect in predicting SMI. Suicidal ideation in the 12 months before interview was assessed with a question that asked respondents whether they had seriously thought about killing themselves in the past year.

Stress buffers—Functional social support^[58] was assessed with one question that asked respondents about the number of people in their county-parish who they could speak to about their private feelings without embarrassment. Social competence was assessed with a 12-item scale^[59] rating such abilities as staying calm in a crisis, getting along with people, being persuasive, staying out of trouble in dangerous situations, staying in control of emotions when necessary, and keeping a sense of humor in tense situations. The internal consistency reliability (Cronbach's α) of this scale was .87.

Data analysis

Prevalence of grief symptoms associated with each type of loss and co-occurrence of CG with SMI-suicidality were examined with cross-tabulations. Predictive associations of socio-demographics, hurricane-related stressors, and pre-hurricane history of psychopathology with CG were examined with logistic regression analysis. Logistic regression coefficients and their standard errors were exponentiated to create odds-ratios (ORs) and 95% confidence intervals (95% CIs). The Taylor series linearization method was used to calculate design-based significance tests. Statistical significance was consistently evaluated using two-sided .05-level tests.

RESULTS

The estimated prevalence of complicated grief

The estimated prevalence of CG in the CAG was 15.3%, representing 26.1% of respondents with a significant loss. (Table 1) The majority of CG (73%) was rated mild, compared to

14.5% moderate and 12.5% severe. Only 4.1% of respondents in the total sample (7.0% of those with a significant loss) were estimated to have moderate (2.2%) or severe (1.9%) CG.

The distribution of grief by type of loss

Only 3.7% of respondents reported that death of a loved one was their most significant hurricane-related loss. Other types of loss were more common, with 58.5% of respondents reporting some type of loss rated 3+ on the 0–10 loss severity scale. (Table 2) The most commonly reported losses were tangible losses (29.0%), interpersonal losses (9.5%), intangible losses (8.1%), and work-financial losses (4.2%). The distribution across the five substantive loss categories did not differ significantly among respondents interviewed 5–6, 7–12, and 13–19 months after the hurricane ($\chi^2_8 = 4.2, p = .84$).

Bereavement had the highest conditional probability of grief symptoms (68.9%) and CG (18.9%). (Table 3, Part I) Grief symptoms were less prevalent among respondents whose main loss was interpersonal (52.5%) or tangible (50.5%). Moderate-severe CG was also lower in these subsamples (10.6% interpersonal, 7.6% tangible). Moderate-severe CG was least common among respondents whose main loss was intangible (1.3%) or work-financial (1.1%). Differences in prevalence of CG by type of most significant loss did not differ among respondents in the three survey waves (i.e., interviewed 5–6, 7–12, and 13–19 months post-hurricane; $\chi^2_8 = 10.3, p = .25$ total CG; $\chi^2_8 = 13.8, p = .09$ moderate-severe CG).

Despite the much higher risk of CG among respondents whose most significant loss was bereavement than other types of loss, only 16.5% of all moderate-severe CG was associated with bereavement. (Table 3, Part II) This proportion did not vary markedly depending on whether respondents were interviewed 5–6 (15.8%), 7–12 (23.2%), or 13–19 (14.2%) months after the hurricane. The highest proportions of moderate-severe CG were associated with tangible (52.9%) and interpersonal (24.0%) losses.

Co-occurrence of complicated grief with Serious Mental Illness and suicidal ideation

The vast majority of respondents with severe CG (83.9%) met criteria for either 30-day SMI or 12-month suicidal ideation. (Table 4) Odds-ratios (ORs) between severe CG (compared to no hurricane-related loss) and these outcomes are in the range 24.0–97.5. Prevalence of SMI or suicidal ideation were considerably lower among respondents with moderate (41.7%) or mild (38.3%) CG (ORs = 7.8–13.3). Prevalence of SMI or suicidal ideation were lower still among respondents with no-minimal or sub-threshold grief symptoms (21.4%), although still with elevated ORs (5.1–6.9) compared to people who had no hurricane-related loss. Prevalence of SMI or suicidal ideation among respondents who had no hurricane-related loss, finally, was 5.1%.

Predictors of complicated grief

We examined associations of socio-demographics, hurricane-related stressors, pre-hurricane history of psychopathology, and post-hurricane resiliency factors with moderate-severe CG among respondents who experienced hurricane-related loss. None of the socio-demographic variables was a significant predictor after controlling for type of loss. This is striking given that socio-demographics typically are associated with more general measures of anxiety and mood disorders in community epidemiological surveys.^[60,61] Access to health insurance, an indirect indicator of socio-economic status, was also unrelated to moderate-severe CG, as was residential location prior to the hurricane (the New Orleans Metropolitan Area vs. the remainder of the areas affected by the hurricane).

Two presumed stress buffers, social support and social competence, were also insignificant predictors of moderate-severe CG. However, the remaining two predictors, number of hurricane-related stressors reported by respondents and pre-hurricane history of psychopathology, were both significant. Number of hurricane-related stressors were coded in the range 1–10 (with 1 being the lowest score rather than 0 because all respondents in the loss subsample reported exposure to at least one hurricane-related stressor), where a score of 10 represents reporting exposure to 10 or more of the 30 stressors assessed in the survey. (Scores were truncated at 10 because only a very small number of respondents reported exposure to more than 10 of the 30 stressors.) The 1.7 OR associated with hurricane-related stressors consequently represents the relative-odds of moderate-severe CG associated with an increase of one stressor. More detailed analyses (results available on request) showed that the implicit assumption of a linear association between number of stressors and log-odds of moderate-severe CG is consistent with the data; that is, that the OR of having been exposed to 3 stressors versus 1 is roughly equal to $1.7^2 = 2.9$; of having been exposed to 4 stressors versus 1 equal to $1.7^3 = 4.9$, etc. Further analysis also supported the model assumption that types of stressors could be considered equivalent for purposes of predicting this outcome.

Pre-hurricane history of psychopathology was coded as a 0–7 count of number of prior lifetime mental disorders assessed in the survey. The OR of 1.2 associated with this predictor consequently represents the association of an increase in one point on this scale. More detailed analyses (results available on request) showed that the implicit assumption of a linear association between number of disorders and log-odds of CG is consistent with the data; that is, that the OR of history of two prior mental disorders vs. 0 is roughly $1.2^2 = 1.4$, of having three prior disorders vs. 0 is roughly $1.2^3 = 1.7$, etc. Further analysis also supported the model assumption that types of disorders are not significant predictors of CG once number of disorders is controlled. This means that the vulnerability associated with history of psychopathology is relatively general rather than linked to any particular subset of disorders.

Decomposition showed that the ORs of the predictors taken as a set are significantly different across subsamples defined by type of loss ($\chi^2_{28} = 47.9$, $p = .011$). However, the only individual predictor for which the ORs are significantly different across subsamples is social competence $\chi^2_2 = 6.4$, $p = .041$). Social competence (standardized to a mean of 0 and variance of 1) has a statistically significant 0.3 OR predicting moderate-severe CG associated with interpersonal loss, but is not significant in predicting moderate-severe CG associated with other types of loss (0.8–0.9). Most other significant specifications involved predictors only of CG associated with bereavement, including elevated ORs associated with Non-White race (6.9), low education (8.6), and social support (2.3). However, caution is needed in interpreting these specifications, as none of the ORs differs significantly from the non-significant ORs for these predictors in the total sample ($\chi^2_2 = 2.2$ – 3.5 , $p = .17$ – $.34$), raising the possibility that significant subsample associations might be due to chance fluctuations in the large number of subsample replications. The only other noteworthy specification is that history of psychopathology does not predict CG associated with bereavement, while it does predict CG associated with other types of loss.

DISCUSSION

The CAG is one of the largest disaster-related surveys that screened for CG and the first to investigate CG associated with non-bereavement disaster-related losses. More than half of respondents reported a disaster-related loss, with types similar to those described after other disasters.^[15,62] More than one-fourth of respondents with a significant loss reported at least some grief symptoms, with moderate-severe CG reported by 3.9% of respondents. Although

CG was significantly associated with SMI and suicidal ideation, the majority of respondents with mild or moderate CG did not have SMI.

CG was most prevalent following bereavement (conditional prevalence of moderate-severe CG of 18.9% compared to 1.1– 10.6% for other losses). These differences were quite consistent across subsamples of respondents that differed in length of time between the hurricane and the time of baseline interview.

Prevalence estimates of CG in other post-disaster studies that assessed CG related to bereavement^[18–20] and other studies of death of a loved one due to a variety of causes^[49,63] vary widely (10–76%). The 18.9% CAG estimate is at the lower end of this range. Caution is needed in interpreting this comparison, though, as each study so far has used a different rating instrument and the CAG estimate was based on a very short screening measure. No other study asked respondents to rate a range of losses and to identify which was most severe.

We are unaware of any previous disaster study that estimated CG associated with non-bereavement loss. Interestingly, because of the comparatively low prevalence of bereavement, other types of loss accounted for the vast majority of CG (83.5%). Property loss was the most common cause of CG (accounting for 52.9% of all CG cases), with interpersonal losses other than death accounting for an additional 24.0%. However, as a result of the high conditional risk of CG among respondents with bereavement, the proportion of CG due to bereavement (16.5%) was a considerably higher proportion than one would expect by chance given that only 6.4% of all respondents who reported a loss said that bereavement was their most significant loss.

The finding of high co-occurrence of CG with both mood-anxiety disorders^[27,64,65] and suicidal ideation^[66–69] is consistent with previous research, and was particularly common (over 80%) among individuals with severe CG. In addition, we found that exposure to hurricane-related stressors was strongly related to CG. This, too, is consistent with previous research.^[18,64] However, our finding that CG from non-bereavement loss was largely unrelated to socio-demographics is inconsistent with evidence from previous studies that bereavement-related CG is generally more common among women, minorities, the unmarried, and people with socioeconomic disadvantage.^[18,25,64] This failure to find strong socio-demographic correlates of CG is part of a larger pattern in the CAG for socio-demographics to be much less strongly related either to trauma exposure or to psychopathology (PTSD or SMI) than in other natural disaster samples.^[41,43] We suggested in a previous report that these weak associations are due to the enormity of the devastation caused by Katrina, which overwhelmed the protective effects typically provided by socio-demographic advantage, leading to a wider distribution of psychopathological reactions than in more typical natural disasters.^[42]

Another CAG finding consistent with previous research is that pre-hurricane history of psychopathology strongly predicted clinically significant CG.^[27,64,70] The finding that *number* of rather than *type* of prior disorders predicted CG is consistent with accumulating evidence that CG is a unique syndrome, not best described as a form of depression or PTSD as many have done,^[1–16] although it is important to be clear that this finding is certainly not definitive in arguing that CG is a unique syndrome. Indeed, one of the weakest aspects of this sample is that it did not include a comprehensive assessment of other disorders with which CG might be confounded.

A series of specifications showed that low education, minority race-ethnic status and social support predicted bereavement-related CG but not other CG, while pre-hurricane history of psychopathology and social competence predicted only non-bereavement-related CG. The

stability of these specifications is uncertain and requires replication in independent datasets. The possibility of specificity, paired with the high prevalence of grief symptoms among those with non-bereavement losses, points to the importance of future studies examining patterns and predictors of grief among individuals who experienced losses other than death of a loved one.

Several observations can be made about these specificities. The finding that markers of disadvantaged social status (minority race-ethnic status, low education) predicted only bereavement-related CG might be taken to suggest that social ties are especially important for people in socially disadvantaged than advantaged positions. Evidence consistent with such a specification exists in the social networks literature.^[71,72] The finding that social support predicts *increased* risk of CG, but only when the CG is related to bereavement, might indicate that social support is a marker of the magnitude of loss rather than a true vulnerability factor. Studies of social support in older bereaved samples indicate an association between greater emotional loneliness and instrumental social support.^[73,74] However, it is unclear how this relates to CG. We are unaware of previous studies that examined effects of social competence on CG. Our finding that social competence does not protect against CG due to non-bereavement loss raises the possibility that protective effects of social competence might be specific to interpersonal losses. All these specifications need to be replicated in other datasets, though, before they are considered reliable.

A number of study limitations are important to note. First, the CAG excluded people who we could not trace as well as those not reachable by telephone, which likely resulted in the under-representation of individuals with the greatest exposure to hurricane-related stressors and, potentially, the highest rates of CG. Second, CG was assessed using a brief symptom scale that included only a subset of the symptoms now recommended for assessment of CG. In addition, symptoms were assessed 5–19 months after the hurricane even though the DSM 5 workgroup suggested a minimum duration of at least 12 months for a diagnosis of bereavement-related adjustment disorder.^[31] Prevalence estimates of CG and results regarding associations should be considered only provisional. It is noteworthy, though, that results regarding correlates of CG did not vary significantly as a function of time between the hurricane and the survey, which means that the patterns reported here are broadly consistent whether CG is defined with durations as short as 4–5 months (i.e., symptoms in the past month among respondents interviewed 5–6 months after the hurricane, 6–11 months, or 12+ months). Third, although the screening scales of co-occurring mental disorders used here have been validated,^[54,56] screening scales are inherently less precise than comprehensive diagnostic interviews, which undoubtedly led to at least some misclassification of respondents. Fourth, it would have been valuable if the survey had included a comprehensive assessment of other disorders that might be comorbid with CG, allowing us to investigate whether or not unique associations could be found with CG after controlling for those other disorders. Fifth, it is difficult to interpret results for the 8.1% of respondents who said that they had a most significant hurricane-related loss that was intangible (e.g., quality of life, sense of well-being, control, security, way of life), as the kinds of experiences included in the description of intangible losses overlap considerably with the symptoms of CG. It is noteworthy, though, that prevalence of clinically significant CG was quite low in this sub-sample (1.3%), minimizing the impact of this problem. These limitations need to be corrected in future studies.

Despite these limitations, the results reported here suggest that CG is associated with losses other than death, that non-death grief might make up a large proportion of CG after a natural disaster, and that the predictors of CG might differ depending on type of loss, although replication of these results in a study that assesses a wide range of other DSM disorders is needed to determine the extent to which these patterns hold up after controlling for all other

relevant disorders. A practical implication of the results for disaster response involves the fact that post-disaster interventions for grief have been developed, although not systematically tested,^[75] and could be applied if clinicians increased their recognition of potentially problematic post-disaster grief reactions. Previous studies indicate that grief-focused therapies out-perform more conventional therapies in ameliorating CG symptoms,^[28–30] but this work has not targeted disaster-bereaved individuals and has not included losses other than death. Data reported here suggest that the magnitude and heterogeneity of the problem of post-disaster CG are large enough to warrant systematic investigation of these possibilities.

Acknowledgments

Funding/Support: This study is supported by NIH Research Grants R01 MH070884-01A2 and R01 MH081832 from the US Department of Health and Human Services, National Institutes of Health (NIH), the Office of the Assistant Secretary of Planning and Evaluation, the Federal Emergency Management Agency, and the Administration for Children and Families, and by NIH Research Grants R01MH070741 and R01MH06078 from the National Institute of Mental Health (NIMH).

Role of the Sponsor: The funding agencies had no role in the design and conduct of the study; in the collection, analysis, and interpretation of the data; or in the preparation, review, or approval of the manuscript.

References

1. Chou FH, Wu HC, Chou P, et al. Epidemiologic psychiatric studies on post-disaster impact among Chi-Chi earthquake survivors in Yu-Chi, Taiwan. *Psychiatry Clin Neurosci*. 2007; 61:370–378. [PubMed: 17610661]
2. DiGrande L, Neria Y, Brackbill RM, et al. Long-term posttraumatic stress symptoms among 3,271 civilian survivors of the September 11, 2001, terrorist attacks on the World Trade Center. *Am J Epidemiol*. 2011; 173:271–281. [PubMed: 21190987]
3. Freedy JR, Saladin ME, Kilpatrick DG, et al. Understanding acute psychological distress following natural disaster. *J Trauma Stress*. 1994; 7:257–273. [PubMed: 8012746]
4. Hobfoll SE, Tracy M, Galea S. The impact of resource loss and traumatic growth on probable PTSD and depression following terrorist attacks. *J Trauma Stress*. 2006; 19:867–878. [PubMed: 17195971]
5. Ironson G, Wynings C, Schneiderman N, et al. Posttraumatic stress symptoms, intrusive thoughts, loss, and immune function after Hurricane Andrew. *Psychosom Med*. 1997; 59:128–141. [PubMed: 9088048]
6. Johannesson KB, Lundin T, Frojd T, et al. Tsunami-exposed tourist survivors: signs of recovery in a 3-year perspective. *J Nerv Ment Dis*. 2011; 199:162–169. [PubMed: 21346486]
7. Kun P, Chen X, Han S, et al. Prevalence of post-traumatic stress disorder in Sichuan Province, China after the 2008 Wenchuan earthquake. *Public Health*. 2009; 123:703–707. [PubMed: 19892379]
8. Kun P, Han S, Chen X, et al. Prevalence and risk factors for posttraumatic stress disorder: a cross-sectional study among survivors of the Wenchuan 2008 earthquake in China. *Depress Anxiety*. 2009; 26:1134–1140. [PubMed: 19842170]
9. Kuo CJ, Tang HS, Tsay CJ, et al. Prevalence of psychiatric disorders among bereaved survivors of a disastrous earthquake in Taiwan. *Psychiatr Serv*. 2003; 54:249–251. [PubMed: 12556609]
10. Norris FH, Friedman MJ, Watson PJ. 60,000 disaster victims speak: Part II. Summary and implications of the disaster mental health research. *Psychiatry*. 2002; 65:240–260. [PubMed: 12405080]
11. Norris FH, Friedman MJ, Watson PJ, et al. 60,000 disaster victims speak: Part I. An empirical review of the empirical literature, 1981–2001. *Psychiatry*. 2002; 65:207–239. [PubMed: 12405079]
12. Raphael B, Ma H. Mass catastrophe and disaster psychiatry. *Mol Psychiatry*. 2011; 16:247–251. [PubMed: 21331093]

13. Smith BW, Freedy JR. Psychosocial resource loss as a mediator of the effects of flood exposure on psychological distress and physical symptoms. *J Trauma Stress*. 2000; 13:349–357. [PubMed: 10838680]
14. Su CY, Tsai KY, Chou FH, et al. A three-year follow-up study of the psychosocial predictors of delayed and unresolved post-traumatic stress disorder in Taiwan Chi-Chi earthquake survivors. *Psychiatry Clin Neurosci*. 2010; 64:239–248. [PubMed: 20602724]
15. Tang CS. Positive and negative postdisaster psychological adjustment among adult survivors of the Southeast Asian earthquake-tsunami. *J Psychosom Res*. 2006; 61:699–705. [PubMed: 17084149]
16. Wahlstrom L, Michelsen H, Schulman A, et al. Different types of exposure to the 2004 tsunami are associated with different levels of psychological distress and posttraumatic stress. *J Trauma Stress*. 2008; 21:463–470. [PubMed: 18956445]
17. Shear MK, Simon N, Wall M, et al. Complicated grief and related bereavement issues for DSM-5. *Depress Anxiety*. 2011; 28:103–117. [PubMed: 21284063]
18. Ghaffari-Nejad A, Ahmadi-Mousavi M, Gandomkar M, et al. The prevalence of complicated grief among Bam earthquake survivors in Iran. *Arch Iran Med*. 2007; 10:525–528. [PubMed: 17903061]
19. Johannesson KB, Lundin T, Hultman CM, et al. The effect of traumatic bereavement on tsunami-exposed survivors. *J Trauma Stress*. 2009; 22:497–504. [PubMed: 19937645]
20. Kristensen P, Weisaeth L, Heir T. Predictors of Complicated Grief After a Natural Disaster: A Population Study Two Years After the 2004 South-East Asian Tsunami. *Death Stud*. 2010; 34:137–150.
21. Bryant RA, Friedman MJ, Spiegel D, et al. A review of acute stress disorder in DSM-5. *Depress Anxiety*. 2010 (e-publication ahead of print).
22. Shear K, Shair H. Attachment, loss, and complicated grief. *Dev Psychobiol*. 2005; 47:253–267. [PubMed: 16252293]
23. Prigerson HG, Bierhals AJ, Kasl SV, et al. Complicated grief as a disorder distinct from bereavement-related depression and anxiety: a replication study. *Am J Psychiatry*. 1996; 153:1484–1486. [PubMed: 8890686]
24. Prigerson HG, Frank E, Kasl SV, et al. Complicated grief and bereavement-related depression as distinct disorders: preliminary empirical validation in elderly bereaved spouses. *Am J Psychiatry*. 1995; 152:22–30. [PubMed: 7802116]
25. Shear KM, Jackson CT, Essock SM, et al. Screening for complicated grief among Project Liberty service recipients 18 months after September 11, 2001. *Psychiatr Serv*. 2006; 57:1291–1297. [PubMed: 16968758]
26. Horowitz MJ, Siegel B, Holen A, et al. Diagnostic criteria for complicated grief disorder. *Am J Psychiatry*. 1997; 154:904–910. [PubMed: 9210739]
27. Simon NM, Shear KM, Thompson EH, et al. The prevalence and correlates of psychiatric comorbidity in individuals with complicated grief. *Compr Psychiatry*. 2007; 48:395–399. [PubMed: 17707245]
28. Shear K, Frank E, Houck PR, et al. Treatment of complicated grief: a randomized controlled trial. *JAMA*. 2005; 293:2601–2608. [PubMed: 15928281]
29. Shear MK, Frank E, Foa E, et al. Traumatic grief treatment: a pilot study. *Am J Psychiatry*. 2001; 158:1506–1508. [PubMed: 11532739]
30. Boelen PA, de Keijser J, van den Hout MA, et al. Treatment of complicated grief: a comparison between cognitive-behavioral therapy and supportive counseling. *J Consult Clin Psychol*. 2007; 75:277–284. [PubMed: 17469885]
31. American Psychiatric Association. DSM-5 development: Adjustment disorders. 2011. [cited January 27, 2011]; Available from: <http://www.dsm5.org/ProposedRevisions/Pages/proposedrevision.aspx?rid=367>
32. Archer J, Rhodes V. The grief process and job loss: a cross-sectional study. *Br J Psychol*. 1993; 84 (Pt 3):395–410. [PubMed: 8401991]
33. Archer J, Hawes J. Grief and rehousing. *Br J Med Psychol*. 1988; 61 (Pt 4):377–379. [PubMed: 3207639]

34. Persinger MA. Personality changes following brain injury as a grief response to the loss of sense of self: phenomenological themes as indices of local lability and neurocognitive structuring as psychotherapy. *Psychol Rep.* 1993; 72:1059–1068. [PubMed: 8337306]
35. Zinner ES, Ball JD, Stutts ML, et al. Modification and factor-analysis of the Grief Experience Inventory in non-death loss bereavement situations. *OMEGA-J Death Dying.* 1991; 23:143–151.
36. Fulton G, Madden C, Minichiello V. The social construction of anticipatory grief. *Soc Sci Med.* 1996; 43:1349–1358. [PubMed: 8913004]
37. Harvey B. Complications of complicated grief in renal failure. *EDTNA ERCA J.* 2000; 26:36–37. 40. [PubMed: 11011636]
38. Herman D, Felton C, Susser E. Mental health needs in New York state following the September 11th attacks. *J Urban Health.* 2002; 79:322–331. [PubMed: 12200501]
39. Loos C, Bowd A. Caregivers of persons with Alzheimer's disease: some neglected implications of the experience of personal loss and grief. *Death Stud.* 1997; 21:501–514. [PubMed: 10175165]
40. Sanders S, Ott CH, Kelber ST, et al. The experience of high levels of grief in caregivers of persons with Alzheimer's disease and related dementia. *Death Stud.* 2008; 32:495–523. [PubMed: 18958942]
41. Galea S, Brewin CR, Gruber M, et al. Exposure to hurricane-related stressors and mental illness after Hurricane Katrina. *Arch Gen Psychiatry.* 2007; 64:1427–1434. [PubMed: 18056551]
42. Kessler RC, Galea S, Gruber MJ, et al. Trends in mental illness and suicidality after Hurricane Katrina. *Mol Psychiatry.* 2008; 13:374–384. [PubMed: 18180768]
43. Kessler RC, Galea S, Jones RT, et al. Mental illness and suicidality after Hurricane Katrina. *Bull World Health Organ.* 2006; 84:930–939. [PubMed: 17242828]
44. McLaughlin KA, Berglund P, Gruber MJ, et al. Recovery from PTSD following Hurricane Katrina. *Depress Anxiety.* 2011 (e-publication ahead of print).
45. Rosenbaum S. US health policy in the aftermath of Hurricane Katrina. *JAMA.* 2006; 295:437–440. [PubMed: 16434635]
46. Abramson, D.; Garfield, R. *On the Edge: Children and Families Displaced by Hurricanes Katrina and Rita Face a Looming Medical and Mental Health Crisis.* New York, NY: Columbia University Mailman School of Public Health; 2008.
47. Prigerson HG, Maciejewski PK, Reynolds CF 3rd, et al. Inventory of Complicated Grief: a scale to measure maladaptive symptoms of loss. *Psychiatry Res.* 1995; 59:65–79. [PubMed: 8771222]
48. Simon NM, Wall MM, Keshaviah A, et al. Informing the symptom profile of complicated grief. *Depress Anxiety.* 2011; 28:118–126. [PubMed: 21284064]
49. Prigerson HG, Bierhals AJ, Kasl SV, et al. Traumatic grief as a risk factor for mental and physical morbidity. *Am J Psychiatry.* 1997; 154:616–623. [PubMed: 9137115]
50. Prigerson HG, Horowitz MJ, Jacobs SC, et al. Prolonged grief disorder: Psychometric validation of criteria proposed for DSM 5 and ICD-11. *PLoS Med.* 2009; 6:e1000121. [PubMed: 19652695]
51. Endicott, J.; Andreasen, N.; Spitzer, R. *Family History Research Diagnostic Criteria.* New York, NY: Biometrics Research, NY State Psychiatric Institute; 1978.
52. Kendler KS, Silberg JL, Neale MC, et al. The family history method: whose psychiatric history is measured? *Am J Psychiatry.* 1991; 148:1501–1504. [PubMed: 1928463]
53. Nock MK, Holmberg EB, Photos VI, et al. Self-Injurious Thoughts and Behaviors Interview: development, reliability, and validity in an adolescent sample. *Psychol Assess.* 2007; 19:309–317. [PubMed: 17845122]
54. Kessler RC, Barker PR, Colpe LJ, et al. Screening for serious mental illness in the general population. *Arch Gen Psychiatry.* 2003; 60:184–189. [PubMed: 12578436]
55. Furukawa T, Kessler R, Slade T, et al. The performance of the K6 and K10 screening scales for psychological distress in the Australian National Survey of Mental Health and Well-Being. *Psychol Med.* 2002; 33:357–362. [PubMed: 12622315]
56. Kessler RC, Andrews G, Colpe LJ, et al. Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychol Med.* 2002; 32:959–976. [PubMed: 12214795]

57. First, M.; Spitzer, R.; Gibbon, M., et al. Structured Clinical Interview for DSM-IV Axis I Disorders, Research Version. New York, NY: Biometrics Research, New York State Psychiatric Institute; 2002. Non-patient Edition (SCID-I/NP)
58. Cohen S, Wills TA. Stress, social support, and the buffering hypothesis. *Psychol Bull.* 1985; 98:310–357. [PubMed: 3901065]
59. Kessler RC, Avenevoli S, Costello EJ, et al. National comorbidity survey replication adolescent supplement (NCS-A): II. Overview and design. *J Am Acad Child Adolesc Psychiatry.* 2009; 48:380–385. [PubMed: 19242381]
60. Kessler, R.; Aguilar-Gaxiola, S.; Alonso, J., et al. Lifetime prevalence and age of onset distributions of mental disorders in the World Mental Health Survey Initiative. In: Kessler, RC.; Üstün, TB., editors. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders.* New York, NY: Cambridge University Press; 2008. p. 511-521.
61. Kessler, RSA-G.; Alonso, J., et al. Prevalence and severity of mental disorders in the WMH Surveys. In: Kessler, RC.; Üstün, TB., editors. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders.* New York, NY: Cambridge University Press; 2008. p. 534-540.
62. Kohn R, Levav I, Donaire I, et al. Psychological and psychopathological reactions in Honduras following Hurricane Mitch: implications for service planning. *Rev Panam Salud Publica.* 2005; 18:287–295. [PubMed: 16354426]
63. Middleton W, Burnett P, Raphael B, et al. The bereavement response: a cluster analysis. *Br J Psychiatry.* 1996; 169:167–171. [PubMed: 8871792]
64. Neria Y, Gross R, Litz B, et al. Prevalence and psychological correlates of complicated grief among bereaved adults 2.5–3. 5 years after September 11th attacks. *J Trauma Stress.* 2007; 20:251–262. [PubMed: 17597124]
65. Silverman GK, Jacobs SC, Kasl SV, et al. Quality of life impairments associated with diagnostic criteria for traumatic grief. *Psychol Med.* 2000; 30:857–862. [PubMed: 11037094]
66. Latham AE, Prigerson HG. Suicidality and bereavement: complicated grief as psychiatric disorder presenting greatest risk for suicidality. *Suicide Life Threat Behav.* 2004; 34:350–362. [PubMed: 15585457]
67. Simon NM, Pollack MH, Fischmann D, et al. Complicated grief and its correlates in patients with bipolar disorder. *J Clin Psychiatry.* 2005; 66:1105–1110. [PubMed: 16187766]
68. Szanto K, Prigerson H, Houck P, et al. Suicidal ideation in elderly bereaved: the role of complicated grief. *Suicide Life Threat Behav.* 1997; 27:194–207. [PubMed: 9260302]
69. Szanto K, Shear MK, Houck PR, et al. Indirect self-destructive behavior and overt suicidality in patients with complicated grief. *J Clin Psychiatry.* 2006; 67:233–239. [PubMed: 16566618]
70. Melhem NM, Rosales C, Karageorge J, et al. Comorbidity of axis I disorders in patients with traumatic grief. *J Clin Psychiatry.* 2001; 62:884–887. [PubMed: 11775048]
71. Turner RJ, Noh S. Class and psychological vulnerability among women: the significance of social support and personal control. *J Health Soc Behav.* 1983; 24:2–15. [PubMed: 6853996]
72. Whyte, W. *Street Corner Society: The Social Structure of an Italian Slum.* Chicago, IL: University of Chicago Press; 1981.
73. Bisconti TL, Bergeman CS, Boker SM. Social support as a predictor of variability: an examination of the adjustment trajectories of recent widows. *Psychol Aging.* 2006; 21:590–599. [PubMed: 16953720]
74. van Baarsen B. Theories on coping with loss: the impact of social support and self-esteem on adjustment to emotional and social loneliness following a partner's death in later life. *J Gerontol B Psychol Sci Soc Sci.* 2002; 57:S33–42. [PubMed: 11773231]
75. Walsh F. Traumatic loss and major disasters: strengthening family and community resilience. *Fam Process.* 2007; 46:207–227. [PubMed: 17593886]

Table 1

Estimated prevalence of grief among adults exposed to Hurricane Katrina (n = 3,088)

	%	(se)
No loss ¹	41.5	(1.5)
No-minimal grief ²	29.4	(1.3)
Sub-threshold grief ²	13.7	(1.0)
Complicated Grief (CG) ²		
Mild	11.1	(0.9)
Moderate ²	2.2	(0.4)
Severe ²	1.9	(0.5)
Total CG	15.3	(1.1)

¹ Scores in the range 0–2 on the 0–10 scale assessing severity of hurricane-related loss. Grief reactions were assessed only among respondents with scores in the range 3–10 on this scale.

² The grief scale was scored in the range 0–16. Scale ranges were defined as 0–4 *no-minimal grief*, 5–7 *sub-threshold grieves*, 8–12 *mild CG*, 13–14 *moderate CG*, and 15–16 *severe CG*.

Table 2Distribution of self-reported most significant hurricane-related losses¹

	%	(se)	(n)
No Loss	41.5	(1.5)	(1,021)
Tangible	29.0	(1.3)	(984)
Interpersonal	9.5	(0.8)	(388)
Intangible	8.1	(0.7)	(330)
Work or financial	4.2	(0.6)	(120)
Death of a loved one	3.7	(0.5)	(160)
No answer/unintelligible	2.7	(0.6)	(51)
Missing	1.2	(0.4)	(34)
Total			(3,088)

¹ Respondents who reported hurricane-related losses in the range 3–10 on the 0–10 severity-of-loss scale were asked to describe their most significant loss. Open-ended responses were coded into the categories reported in this table. Tangible losses include such things as loss of a home, possessions, or memorabilia. Interpersonal losses (other than death of a loved one) include separations from family or friends as well as decreases in quality of relationships with family or friends. Intangible losses include such things as loss of quality of life, sense of well-being, control, security, or way of life. Work/financial losses include any mention of job, money, finance, income, or business losses.

Table 3

Estimated prevalence and severity of complicated grief (CG) according to type of loss¹

	No-minimal grief (0-4)		Sub-threshold grief (5-7)		Mild CG (8-12)		Moderate CG (13-14)		Severe CG (15-16)		Moderate/Severe CG (13-16)	
	%	(se)	%	(se)	%	(se)	%	(se)	%	(se)	%	(se)
I. Percentage by rows												
Tangible Losses	49.5	(2.7)	20.8	(2.2)	22.0	(2.4)	4.9	(1.1)	2.7	(1.1)	7.6	(1.5)
Interpersonal Losses	47.5	(4.6)	28.4	(4.2)	13.5	(2.8)	2.8	(1.2)	7.7	(3.2)	10.6	(3.3)
Intangible Losses	57.4	(4.4)	26.9	(4.3)	14.4	(2.7)	0.9	(0.6)	0.4	(0.3)	1.3	(0.6)
Work/Financial Losses	64.4	(7.0)	15.3	(4.7)	19.2	(6.0)	1.1	(0.8)	0.0	(0.0)	1.1	(0.8)
Death of a Loved One	31.1	(6.4)	22.0	(5.3)	28.0	(5.7)	11.4	(4.9)	7.5	(2.9)	18.9	(5.4)
Unintelligible	60.3	(10.9)	31.5	(10.9)	4.3	(2.2)	0.3	(0.3)	3.6	(3.2)	4.0	(3.2)
Missing	31.7	(13.0)	40.4	(13.7)	26.7	(15.2)	0.0	(0.0)	1.2	(1.2)	1.2	(1.2)
II. Percentage by columns												
Tangible Losses	48.8	(2.6)	44.0	(4.0)	57.4	(4.2)	63.9	(8.6)	40.2	(12.6)	52.9	(7.5)
Interpersonal Losses	15.4	(1.9)	19.7	(3.1)	11.6	(2.5)	11.9	(5.1)	38.0	(12.5)	24.0	(6.9)
Intangible Losses	15.8	(1.7)	15.8	(2.8)	10.5	(2.0)	3.2	(2.1)	1.8	(1.4)	2.5	(1.3)
Work/Financial Losses	9.2	(1.7)	4.7	(1.5)	7.3	(2.4)	2.1	(1.6)	0.0	(0.0)	1.1	(0.8)
Death of a Loved One	3.9	(1.0)	5.9	(1.5)	9.2	(2.1)	18.6	(7.7)	14.1	(5.9)	16.5	(5.0)
Unintelligible	5.6	(1.5)	6.3	(2.6)	1.1	(0.5)	0.4	(0.4)	5.1	(4.5)	2.6	(2.1)
Missing	1.3	(0.6)	3.6	(1.4)	3.0	(2.0)	0.0	(0.0)	0.8	(0.8)	0.4	(0.4)
(n)	(1,132)	(461)	(358)	(77)	(39)	(116)						

¹The 2,183 respondents in this table consist of all those who reported hurricane-related losses in the range 3-10 on the 0-10 severity-of-loss scale. The sub-sample n's do not correspond to the proportions in Table 1-2 due to the fact that those proportions are based on weighted data, while the n's reported in the current table are unweighted.

Table 4

Co-occurrence of complicated grief (CG) with 30-day serious mental illness (SMI) and 12-month suicidal ideation (n = 3,088)^f

	30-Day SMI ^f			12-Month suicidal ideation			SMI or suicidal ideation				
	%	(se)	OR (95% CI)	%	(se)	OR (95% CI)	%	(se)	OR (95% CI)		
No hurricane-related loss	4.2	(0.7)	1.0	--	1.0	(0.4)	1.0	(0.8)	1.0	--	
No-minimal/sub- threshold grief	18.7	(3.1)	5.3*	(3.0-9.1)	6.5	(2.2)	6.9*	(2.4-19.9)	21.4	(3.3)	5.1* (3.0-8.5)
Mild CG	35.6	(4.2)	12.6*	(7.5-21.1)	7.3	(2.4)	7.8*	(2.8-22.1)	38.3	(4.3)	11.6* (7.1-18.9)
Moderate CG	39.8	(8.3)	15.1*	(7.0-32.5)	9.5	(4.7)	10.4*	(2.8-38.8)	41.7	(8.5)	13.3* (6.2-28.4)
Severe CG	79.3	(9.4)	87.2*	(27.3-278.9)	19.5	(10.2)	24.0*	(5.5-105.6)	83.9	(8.7)	97.5* (26.7-356.3)
Any CG	41.7	(3.8)	16.4*	(10.2-26.3)	9.2	(2.3)	10.0*	(3.9-25.9)	44.6	(3.8)	15.0* (9.5-23.6)
Moderate-severe CG	58.1	(7.5)	31.6*	(15.7-63.7)	14.1	(5.4)	16.3*	(5.1-52.6)	61.3	(7.4)	29.5* (14.7-59.3)

^f χ^2_4 tests for the association between level of grief (no loss, no-minimal/sub-threshold grief, mild CG, moderate CG, severe CG) and each dichotomous outcome are: SMI $\chi^2_4 = 135.7$, $p < 0.001$, suicidal ideation $\chi^2_4 = 26.3$, $p < 0.001$, SMI or suicidal ideation $\chi^2_4 = 135.7$, $p < 0.001$

Table 5

Multivariate associations of socio-demographics, hurricane-related stressors, pre-hurricane history of psychopathology, and post-hurricane resiliency factors with moderate-severe complicated grief (CG) among respondents with any hurricane-related loss and separately among respondents with the hurricane-related losses most strongly associated with CG¹

	Any loss		Death of a loved one		Tangible Losses		Interpersonal losses	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Age								
18-39	0.5	(0.2-1.5)	3.2	(0.5-21.2)	0.3	(0.1-1.1)	0.2	(0.0-2.6)
40-59	0.8	(0.3-2.5)	2.5	(0.4-16.6)	0.9	(0.3-2.9)	0.1	(0.0-1.5)
60+	1.0	--	1.0	--	1.0	--	1.0	--
χ^2_2	2.5		1.5		4.5		2.8	
Sex								
Male	0.7	(0.3-1.3)	0.4	(0.1-1.4)	0.6	(0.2-1.5)	1.5	(0.4-5.3)
Female	1.0	--	1.0	--	1.0	--	1.0	--
χ^2_1	1.4		2.3		1.3		0.3	
Race/ethnicity								
Non-white	1.8	(0.9-3.5)	6.9*	(1.9-25.0)	1.7	(0.7-4.3)	1.4	(0.3-6.6)
White	1.0	--	1.0	--	1.0	--	1.0	--
χ^2_1	2.8		8.7*		1.2		0.2	
Income								
Low/low-middle	1.9	(0.9-4.0)	1.5	(0.4-6.0)	2.1	(0.6-7.1)	3.6	(0.6-23.2)
Middle-high	1.0	--	1.0	--	1.0	--	1.0	--
χ^2_1	3.2		0.4		1.5		1.9	
Health insurance								
None	0.9	(0.5-1.9)	0.4	(0.1-1.3)	0.8	(0.2-2.7)	0.6	(0.1-2.9)
Any	1.0	--	1.0	--	1.0	--	1.0	--
χ^2_1	0.0		2.6		0.1		0.4	
Education (in years)								
0-12	1.5	(0.8-2.9)	6.6*	(1.4-31.2)	1.3	(0.5-3.9)	2.1	(0.6-7.8)
13+	1.0	--	1.0	--	1.0	--	1.0	--

	Any loss		Death of a loved one		Tangible Losses		Interpersonal losses	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
χ^2_1	1.4		5.7*		0.3		1.3	
Pre-hurricane residence								
New Orleans metro	1.6	(0.8–3.1)	4.8	(0.6–36.2)	1.3	(0.4–3.6)	1.0	(0.3–3.9)
All others	1.0	--	1.0	--	1.0	--	1.0	--
χ^2_1	2.0		2.4		0.2		0.0	
Marriage status								
Married	0.7	(0.3–1.6)	0.5	(0.1–2.6)	0.6	(0.2–2.2)	1.5	(0.2–10.4)
Previously married	0.6	(0.2–1.4)	2.5	(0.5–12.4)	0.4	(0.1–1.3)	0.5	(0.0–7.6)
Never married	1.0	--	1.0	--	1.0	--	1.0	--
χ^2_2	1.5		3.6		2.7		0.8	
Hurricane-related stressors ²								
	1.7*	(1.5–2.1)	1.4*	(1.0–1.8)	1.9*	(1.4–2.5)	1.9*	(1.2–2.9)
χ^2_1	45.5*		4.3*		20.0*		7.3*	
Social support ³								
	1.2	(0.9–1.7)	2.3*	(1.1–4.8)	1.2	(0.8–2.0)	1.1	(0.4–3.1)
χ^2_1	1.7		5.0*		0.9		0.0	
Social competence ³								
	0.7	(0.6–1.0)	0.8	(0.5–1.3)	0.9	(0.6–1.4)	0.3*	(0.2–0.6)
χ^2_1	3.7		1.1		0.2		11.8*	
Pre-hurricane history of psychopathology ⁴								
	1.2*	(1.0–1.4)	1.2	(0.8–1.7)	1.2*	(1.0–1.5)	1.5*	(1.1–2.1)
χ^2_1	6.0*		1.1		4.5*		6.2*	
(n)	(2,183)		(160)		(984)		(388)	

* Significant at the .05 level, two-sided test

¹ Multivariate logistic regression models predicting moderate-severe complicated grief among respondents who reported hurricane-related losses in the range 3–10 on the 0–10 severity-of-loss scale.

²The hurricane-related stressors scale is a count of number of hurricane-related stressors out of the 30 assessed. The scale is scored in the range 1–10 in this sub-sample. The lowest score is 1 because all respondents with significant loss reported at least one hurricane-related stressor. The highest score is 10 because the small number of respondents who reported more than 10 hurricane-related stressors were coded 10 due to the rarity of higher exposures.

³The social support and social competence scales are standardized to have a mean of 0.0 and a variance of 1.0 in the total sample.

⁴Pre-hurricane history of psychopathology is coded in the range 0–7 and represents the number of lifetime disorders the respondent reported having prior to the hurricane. These seven include major depressive episode, panic disorder, generalized anxiety disorder, post-traumatic stress disorder, intermittent explosive disorder, substance abuse with or without dependence, and suicide ideation.